

The group of automorphisms of semigroup of endomorphisms of free commutative and free associative algebras

A. Berzins

University of Latvia

e-mail: aberzins@latnet.lv

August 2004

AMS Mathematics Subject Classification: 14A99, 14P05

Keywords: commutative algebra, associative algebra, automorphism, endomorphism, variety

Abstract

In this paper are described the groups of automorphisms of semigroup $\text{End}(W(X))$, where $W(X)$ is free commutative or free associative algebra.

1 Introduction

The basis of the classical algebraic geometry is the Galois correspondence between P -closed ideals in $P[x_1, x_2, \dots, x_n]$ and algebraic sets in the affine space P^n . Having noted that a point $a = (a_1, a_2, \dots, a_n)$ uniquely determines a homomorphism $x \in \text{Hom}(P[X])$ ($s(x_i) = a_i$), B. Plotkin in [11-13] makes foundation of the universal algebraic geometry, i.e., algebraic geometry for arbitrary variety Θ of universal algebras. Let us fix a variety Θ and an algebra $H \in \Theta$. Let $W = W(X)$ be a free algebra in Θ with finite set of generators. The set $\text{Hom}(W, H)$ we consider as affine space of points over

H . Then arbitrary congruence T in W determines the set of points in affine space $\text{Hom}(W, H)$:

$$T'_H = A = \{\mu \in \text{Hom} | T \subset \text{Ker} \mu\}.$$

A set of points $A \subset \text{Hom}(W, H)$ determines congruence of W

$$A' = A'_W = T = \bigcap_{\mu \in A} \text{Ker} \mu.$$

We call the set of points A such that $A = T'$ for some T an algebraic set in $\text{Hom}(W, H)$. A relation T with $T = A'$ for some T is a congruence in W . We call such a congruence an H -closed one.

It was found [3, 4, 11-14] that many problems in the universal algebraic geometry, such as geometric equivalence, geometric similarity, isomorphism and equivalence of categories of algebraic sets and varieties depend on structure of $\text{Aut}(\Theta^0)$ and $\text{Aut}(\text{End}(W))$, where Θ^0 is the category of algebras $W = W(X)$ with the finite X that are free in Θ , and W is a free algebra in Θ . Structure of $\text{Aut}(\text{Com} - P)^0$ (i.e, the classical case) was described in [2]. Here was introduced variant of the concept of quasiinner automorphism - basic concept for description of $\text{Aut}(\text{End}(W))$ and $\text{Aut}(\Theta^0)$. Later groups $\text{Aut}(\Theta^0)$ were described for categories of free Lie algebras and free associative algebras [1, 5, 8, 9] and for categories of free modules and free Lie modules [5, 6]. Some other definition of quasiinner automorphism was used in mentioned papers. Also are described $\text{Aut}(\Theta^0)$ for varieties of groups, semigroups and some another varieties. For some varieties is described $\text{Aut}(\text{End}(W))$, but these problems are more difficult. In particular, this problem was not solved even in the classical case of free commutative algebras, i.e. the structure of $\text{Aut}(\text{End} P[x_1, \dots, x_n])$ is not describe if $n > 2$. In all solved cases groups $\text{Aut}(\text{End}(W))$ are generated by semi-inner automorphisms and mirror automorphism. In [3] this problem in classical case was solved, if the field P is algebraically closed.

I want to note that the proof in [3] does not use the description of the group $\text{Aut}(P[X])$. It is very important, because structure of the group $\text{Aut}(P[x_1, \dots, x_n])$ is not described, when $n > 2$, but counterexamples show that structures of $\text{Aut}(P[x_1, \dots, x_n])$ and $\text{Aut}(P[x, y])$ are principally different. Now in the presented paper the group $\text{Aut}(\text{End}(W_n))$ is described for commutative and associative algebras over arbitrary infinite field.

2 Definitions

Let us recall some definitions for variety of commutative algebras $\text{Com} - P$ (see [4,13] for general case).

Definition 2.1 *Let $P[X] = P[x_1, \dots, x_n]$ be a free commutative algebra over a field P with a finite set of generators and $\tau \in \text{Aut}(\text{End}(P[X]))$. It is known [4] that there exists a bijection $\mu : P[X] \mapsto P[X]$ such that for every $s \in \text{End}(P[X])$*

$$s^\tau = \mu s \mu^{-1}.$$

Such representation of τ is called a representation of τ as quasiinner automorphism generated by μ .

Of course, arbitrary set of substitutions does not generate an automorphism, But there are two kinds of bijections μ in commutative case and three in the associative case that generate an automorphism.

1. $\mu = \bar{\alpha}$, where $\bar{\alpha}$ is the natural extension of an automorphism $\alpha \in \text{Aut}(P[X])$ on W , i.e. $\bar{\alpha}(\sum a_k u_k) = \sum \alpha(a_k) u_k$. We shall write $\bar{\alpha} = \alpha$ and call it "automorphism of field".

2. $\mu = \eta \in \text{Aut}(P[X])$.

Definition 2.2 *The automorphism τ generated by an automorphism $\mu \in \text{Aut}(P[X])$ is called inner automorphism.*

Definition 2.3 *Note, that every $\alpha \in \text{Aut}(P)$ belongs to normalizer of subgroup $\text{Aut}(P[X])$ in the group of all bijections of $P[X]$. So, every product of elements of $\text{Aut}(P)$ and $\text{Aut}(P[X])$ may be represented in the form $\mu = \alpha\eta$, $\alpha \in \text{Aut}P$, $\eta \in \text{Aut}(P[X])$. Automorphism $\tau \in \text{Aut}(\text{End}(P[X]))$ generated by μ is called semiinner.*

3. Now we describe the third type of μ for associative algebras. Let $S = S(X)$ be a free semi-group. For every $u = x_{i_1} x_{i_2} \dots x_{i_n}$ in S , take $\bar{u} = x_{i_n} \dots x_{i_2} x_{i_1}$. Then $u \rightarrow \bar{u}$ is an antiautomorphism of the semigroup S .

Definition 2.4 Let $W = W(X)$ be a free associative algebra. For every its element $\omega = \lambda_0 + \lambda_1 u_1 + \dots + \lambda_k u_k$ denote $\beta(\omega) = \overline{\omega} = \lambda_0 + \lambda_1 \overline{u_1} + \dots + \lambda_k \overline{u_k}$. The transition $\beta : W(X) \rightarrow W(X)$ is an antiautomorphism of the algebra W , and it generates the automorphism $\delta \in \text{Aut}(\text{End}(W(X)))$, $s^\delta = \beta s \beta^{-1}$. We call β mirror antiautomorphism of W and δ mirror automorphism of $\text{Aut}(\Theta^0)$.

Here δ is not inner and is not semi-inner, but is quasi-inner. Note that

1. δ belongs to the normalizer of subgroup $\text{Sinn}(\text{End}(W(X)))$ in $\text{Aut}(\text{End}(W(X)))$ that consists of all semi-inner automorphisms;
2. every antiautomorphism of W is a product of the mirror antiautomorphism β and an automorphism of W .

Let now $W(X)$ be a free finitely generated algebra in arbitrary variety Θ .

Definition 2.5 Bijection $\mu : W(X) \mapsto W(X)$ is called central, if for every $s \in \text{End}(W(X))$

$$\mu s = s \mu.$$

Algebra $W(X)$ is called central, if every its central bijection is identical.

Theorem 1 Let $W(X)$ be a free central algebra and μ some bijection of $W(X)$ generating an automorphism $\tau \in \text{Aut}(\text{End}(W(X)))$. Then μ transforms every base of algebra $W(X)$ to a base of algebra.

Proof. [see 4]

For free algebras in a variety Θ of algebras over field P (Com-P, Ass-P, and others) we shall correct the definition of central algebra. Clearly, in every such algebra $W(X)$ we have a bijection $\mu(u) = au + b$, where $a, b \in P$, $a \neq 0$ (linear bijection). This bijection commutate with every endomorphism of algebra $W(X)$ and, obviously, transforms every base of $W(X)$ to a base.

Definition 2.6 Finitely generated algebra $W(X)$ over field P is called almost central, if every its central bijection is linear.

Theorem 2 Free finitely generated commutative, associative are almost central.

Proof. Let μ be a central bijection of $W(X)$, $\mu(x_i) = r_i(x_1, x_2, \dots, x_n) \in W(X)$, and $s \in \text{End}(W(X))$, $s(x_1) = x_1$, $s(x_i) = 0$ for $i = 1$. Then

$$\mu s(x_1) = \mu(x_1) = r_1(x_1, x_2, \dots, x_n),$$

$$s\mu(x_1) = s(r_1(x_1, x_2, \dots, x_n)) = r_1(x_1, 0, \dots, 0) = r(x_1).$$

So $r_1(x_1, x_2, \dots, x_n) = r(x_1)$ is a polynomial of one variable.

For arbitrary $u \in W(X)$ take $s \in \text{End}(W(X))$ $s(x_1) = u$, $s(x_i) = x_i$ for $i > 1$. Then

$$\mu(u) = \mu s(x_1) = s\mu(x_1) = s(r(x_1)) = r(u),$$

and so $\mu(u) = r(u) = a_0 + a_1u + \dots + a_ku^k$. Because μ is bijection, so r is linear, Q.E.D..

Corollary. *From the theorems 1 and 2 we have that in mentioned algebras every bijection μ , which generate an automorphism of $\text{End}(W(X))$ transform every base of algebra $W(X)$ to a base of this algebra.*

3 Linearity of automorphism generating bijection

Let $W = W(X)$ be a free commutative, associative or Lee algebra over the field P with a finite set of generators, and τ be an automorphism of $\text{End}(W(X))$ generated by bijection μ . We call $s \in \text{End}(W(X))$ a constant endomorphism if $\text{Im}(s) = P$. Clearly, if s is constant then s^τ also is constant. Denote the set of all constant endomorphisms by Const .

Let $a \in P$ and s be a constant endomorphism such that $s(x_1) = a$. Then $\mu(a) = \mu(s(x_1)) = s^\tau(\mu(x_1)) \in P$. So a restriction of μ on P is a bijection: $\mu : P \mapsto P$.

Let $\mu(0) = a$ and $\mu(1) = b$. We shall consider the linear bijection $l : W(X) \mapsto W(X)$, $l(u) = cu + d$ such that $l(a) = 0$ and $l(b) = 1$. Since linear bijection generates identical automorphism of the semigroup $\text{End}(W(X))$, so bijections μ and $\mu' = l\mu$ generate equal automorphisms. Note that $\mu'(0) = 0$ and $\mu'(1) = 1$. We denote $\mu = \mu'$ and assume that $\mu(0) = 0$ and $\mu(1) = 1$.

Since μ transforms base of algebra $W(X)$ to base, so the endomorphism $\eta : \eta(x_i) = \mu(x_i)$ is an automorphism of $W(X)$. Now we shall consider the bijection $\mu' = \eta^{-1}\mu$. We have for μ' that $\mu'(x_i) = x_i$.

Now we shall consider the Galois correspondence between ideals in $W(X)$ and algebraic sets in affine space over $W(X)$. We recall that every ideal T in $W(X)$ defines an algebraic set of points in affine space over $W(X)$.

$$T' = T'_{W(X)} = A = \{\eta \in \text{End}(W(X)) | T \subset \text{Ker} \eta\}.$$

A set of points $A \subset \text{End}(W(X))$ determines ideal in $W(X)$:

$$A' = A'_W = \bigcap_{\eta \in A} \text{Ker } \eta.$$

An element $u \in W(X)$ is called basic, if it belongs to some base of algebra $W(X)$.

Lemma 3.1 *If u is an basic element of $W(X)$, then*

$$\langle u \rangle'' = \langle u \rangle$$

Proof. Clearly,

$$A = \langle u_1 \rangle' = \{\eta \in \text{End} W | \eta(u_1) = 0\}$$

and

$$A' = \langle u_1 \rangle'' = \bigcap_{\eta \in A} \text{Ker } \eta = \bigcap_{\eta(u_1)=0} \text{Ker } \eta.$$

Let $f(u_1, u_2, \dots, u_n) \in \langle u_1 \rangle''$ and $\eta : \eta(u_1) = 0, \eta(u_i) = u_i$ for $i > 1$. Then $0 = \eta(f(u_1, u_2, \dots, u_n)) = f(0, u_2, \dots, u_n)$, so $f \in \langle u_1 \rangle$, Q.E.D..

Lemma 3.2 *Let μ be an automorphism generating bijection of W such that $\mu(0) = 0, \mu(1) = 1$ and $\mu(x_i) = x_i$. Then*

$$\langle \mu(\alpha x_1) \rangle = \langle \mu(x_1) \rangle = \langle x_1 \rangle$$

Proof. At first note that for any endomorphism $s \in \text{End}(W)$

$$s(x_1) = 0 \Leftrightarrow s^\tau(x_1) = 0.$$

Really, if $s(x_1) = 0$, then $s^\tau(x_1) = s^\tau \mu(x_1) = \mu s(x_1) = 0$; and if $s^\tau(x_1) = 0$, then $0 = s^\tau(x_1) = s^\tau \mu(x_1) = \mu s(x_1) = 0$, i.e. $s(x_1) = 0$.

Let $\mu(\alpha x_1) = r_1(x_1, \dots, x_n)$. Take $s \in \text{End}(W)$ such that $s^\tau(x_1) = 0$ and $s^\tau(x_i) = x_i$ for $i > 1$. Then $0 = \mu(0) = \mu s(\alpha x_1) = s^\tau \mu(\alpha x_1) = s^\tau r_1(x_1, \dots, x_n) = r_1(0, x_1, \dots, x_n)$, i.e. $\mu(\alpha x_1) \in \langle x_1 \rangle'' = \langle x_1 \rangle$.

Now let us prove, that $x_1 \in \langle \mu(\alpha x_1) \rangle$. Note that

$$\langle \mu(\alpha x_1) \rangle = \langle \mu(\alpha x_1) \rangle'' = \bigcap_{\eta^\tau: \eta^\tau(\mu(\alpha x_1))=0} \text{Ker } \eta^\tau.$$

So we must prove that, if $\eta^\tau(\mu(\alpha x_1)) = 0$ then $\eta^\tau(x_1) = 0$. Really,

$$0 = \eta^\tau(\mu(\alpha x_1)) = \mu \eta(\alpha x_1) = \mu(\alpha \eta(x_1)) \Rightarrow \alpha \eta(x_1) = 0 \Rightarrow \eta^\tau(x_1) = 0.$$

Q.E.D.

Lemma 3.3 *Let μ be an automorphism generating bijection of W such that $\mu(0) = 0$, $\mu(1) = 1$ and $\mu(x_i) = x_i$. Then for every $a \in P$ and $u \in P[X]$*

$$\mu(au) = \mu(a)\mu(u).$$

Proof. Since $\langle \mu(ax_1) \rangle = \langle x_1 \rangle$, so $\mu(ax_1) = bx_1 = b\mu(x_1)$, where $w \in P[X]$. Substitution $s(x_1) = 1$ gives that $b = \mu(a)$, i.e. $\mu(ax_1) = \mu(a)\mu(x_1)$.

In associative case we consider only the case, where the field P is of characteristic 0. I must note, that $\mu(ax_1)$ together with $x_2, \dots, x_n$ is a base of algebra, and so $\mu(ax_1) = bx_1 + f(x_2, \dots, x_n)$ (see [7]). Because $\mu(ax_1) \in \langle x_1 \rangle$, so $\mu(ax_1) = bx_1$.

Lemma 3.4 *With the preceding notation*

$$\mu(u + v) = \mu(u) + \mu(v).$$

Proof. Let $\mu(x_1 + x_2) - \mu(x_1) - \mu(x_2) = f(x_1, \dots, x_n) = g(\mu(x_1), \dots, \mu(x_n))$, then $\mu(u_1 + u_2) - \mu(u_1) - \mu(u_2) = g(\mu(u_1), \dots, \mu(u_n))$. Substitution $s: s(u_1) = u_1, s(u_2) = u_2, s(u_i) = 0$ for $i > 2$ gives identity

$$\mu(u_1 + u_2) - \mu(u_1) - \mu(u_n) = \bar{g}(\mu(u_1), \mu(u_2)).$$

Clearly, $\bar{g}$ is homogenous of degree 1 of u_1 and u_2 . Therefore $\bar{g}(u_1, u_2) = au_1 + bu_2$ and

$$\mu(u + v) = \mu(u) + \mu(v) + a\mu(u) + b\mu(v).$$

Substitutions $u = 0$ and $v = 0$ gives that $a = b = 0$, Q.E.D..

Corollary. Note that restriction μ on P is an automorphism of P and denote it η . Let $\mu' = \eta^{-1}\mu$, then μ' is identical on P . So for $\mu = \mu'$ we have identity

$$\mu(au) = a\mu(u), \tag{1}$$

for every $a \in P$ and $u \in W(X)$.

4 Description of $\text{Aut}(\text{End}(W(X)))$

Let consider at first the commutative case.

Lemma 4.1 *Let μ be a bijection of $P[X]$ generating an automorphism $\tau \in \text{Aut}(\text{End}(P[X]))$ such that μ is identical on P and $\mu(x_i) = x_i$. Then for every $u, v \in P[X]$ it holds that*

$$\mu(u \cdot v) = \mu(u) \cdot \mu(v)$$

Proof. Let s be a constant endomorphism of $P[X]$. Then s^τ also is constant and so

$$\begin{aligned} s^\tau(\mu(x_1x_2) - \mu(x_1)\mu(x_2)) &= s^\tau\mu(x_1x_2) - s^\tau\mu(x_1) \cdot s^\tau\mu(x_2) = \\ \mu s(x_1x_2) - \mu s(x_1) \cdot \mu s(x_2) &= \mu(a_1a_2) - \mu(a_1) \cdot \mu(a_2) = 0, \\ \mu(x_1x_2) - \mu(x_1)\mu(x_2) &\in \bigcap_{s^\tau \in \text{Const}} \text{Ker } s^\tau = 0, \end{aligned}$$

Q.E.D..

Now we can formulate the main result for commutative case.

Theorem 3 *Every automorphism of $\text{End}(P[X])$ is semiinner.*

Proof. It follows from lemma 3.4, lemma 4.1 and identity (1).

Now let consider associative case.

Lemma 4.2 *Let μ be a bijection of $W(X)$ generating an automorphism $\tau \in \text{Aut}(\text{End}(W(X)))$ such that μ' is identical on P and $\mu(x_i) = x_i$. Then for every $u, v \in W(X)$ it holds that*

$$\mu(u \cdot v) = \mu(u) \cdot \mu(v),$$

or for every $u, v \in W(X)$ it holds that

$$\mu(u \cdot v) = \mu(v) \cdot \mu(u),$$

Proof. Let $\mu(x_1x_2) - \mu(x_1)\mu(x_2) = f(x_1, \dots, x_n) = g(\mu(x_1), \dots, \mu(x_n)) = \bar{g}(\mu(x_1), \mu(x_2))$. Note that $\bar{g}$ is 1-homogenous of x_1 and 1-homogenous of x_2 . Following the proof of lemma 3.4 we get identity

$$\mu(uv) = a\mu(u)\mu(v) + b\mu(v)\mu(u).$$

Substitution $v = 1$ gives equality $a + b = 1$. We shall denote $\mu(u) = \bar{u}$ and calculate $\mu(xxy)$.

$$\begin{aligned}\mu(xxy) &= a\bar{x}\bar{x} \cdot \bar{y} + b\bar{y} \cdot \bar{x}\bar{x} = \\ &= a(a+b)\bar{x} \cdot \bar{x} \cdot \bar{y} + b(a+b)\bar{y} \cdot \bar{x} \cdot \bar{x} = \\ &= a\bar{x} \cdot \bar{x} \cdot \bar{y} + b\bar{y} \cdot \bar{x} \cdot \bar{x}, \\ \mu(xxy) &= a\bar{x} \cdot \bar{x}\bar{y} + b\bar{x}\bar{y} \cdot \bar{x} = \\ &= a^2\bar{x} \cdot \bar{x} \cdot \bar{y} + 2ab\bar{x} \cdot \bar{y} \cdot \bar{x} + b^2\bar{y} \cdot \bar{x} \cdot \bar{x}\bar{y}.\end{aligned}$$

So $a = a^2$, $b = b^2$, $ab = 0$, $a + b = 1$ and we have two solutions $a = 1, b = 0$ or $a = 0, b = 1$, Q.E.D..

Now we can formulate the main result for associative algebras.

Theorem 4 *The group $\text{Aut}(\text{End}(W(X)))$ is generated by semiinner automorphisms and mirror automorphism.*

Proof. It follows from lemma 3.4, lemma 4.2 and identity (1).

Acknowledgments

The author is happy to thank professor B. Plotkin for stimulating discussions of the results.

References

1. A. Berzins, *The automorphisms of $\text{End}K[x]$* , Proc. Latvian Acad. Sci., Section B, 2003, vol. 57, no. 3/4, pp. 78-81
2. A. Berzins, *Geometric equivalence of algebras*, Int. J. Alg. Comput., 2001, vol. 11, no. 4, pp. 447-456.

3. A. Berzins, *The group of automorphisms of semigroup $End(P[X])$* , (to appear)
4. A. Berzins, B. Plotkin and E. Plotkin, *Algebraic geometry in Varieties with the Given Algebra of constants*, J. Math. Sci., New York, 2000, vol. 102, no. 3, pp. 4039-4070
5. R. Lipyanski, B. Plotkin *Automorphisms of categories of free modules and free Lie algebras*, ArXiv math.RA/0502212 (10 Feb 2005) pp. 14.
6. Y. Katsov, R. Lipyanski, B. Plotkin, *Automorphisms of categories of free modules, free semimodules and free Lie modules*, (to appear)
7. L. Makar-Limanov *Algebraically closed skew field*, Journal of Algebra, Vol. 93, 1985, pp. 117-135.
8. G. Mashevitzky, *Automorphisms of categories of free associative algebras.*, Preprint.
9. G. Mashevitzky, B. Plotkin, E. Plotkin, *Automorphisms of categories of free Lie algebras.*, Journal of Algebra, Vol. 282, 2004, pp. 490-512.
10. B. Plotkin, *Algebraic logic, varieties of algebras and algebraic varieties*, Proc. Int. Alg. Conf., St. Petersburg, 1995, Walter de Gruyter, New York, London, 1996.
11. B. Plotkin, *Varieties of algebras and algebraic varieties*, Israel J. of Mathematics, 1996, vol. 96, no. 2, pp. 511-522.
12. B. Plotkin, *Varieties of algebras and algebraic varieties. Categories of algebraic varieties*, Sib. Adv. Math., 1997, vol. 7, no. 2, pp. 64-97.
13. B. Plotkin, *Algebras with the same (algebraic) geometry*, Proc. Steklov Inst. Math, Vol. 242, 2003, pp. 165-196.
14. B. Plotkin, G Zhitomirski, *On automorphisms of categories of universal algebras*, ArXiv math.CT/0411408 (18 Nov 2004). pp. 24.